

1. Solve for all values of  $x$  in the interval  $0^\circ \leq x < 360^\circ$  :

$$2 \sin^2 x + 3 \cos x - 3 = 0$$

1.  
  
\_\_\_\_\_

2. Solve for all values of  $x$  in the interval  $0^\circ \leq x < 360^\circ$  :

$$5 \cos^2 x - 6 \sin x = 0$$

2.  
  
\_\_\_\_\_

3. Solve for all values of  $x$  in the interval  $0^\circ \leq x < 360^\circ$  :

$$5 \sec^2 x - 11 \tan x - 4 = 0$$

3.  
  
\_\_\_\_\_

4. Prove the identity:  $\frac{1 + \sec x}{\csc x} = \sin x + \tan x$

4.  
  
\_\_\_\_\_

5. Prove the identity:  $\frac{\sec^2 x - \tan^2 x}{2 \sin^2 x + 2 \cos^2 x} = \frac{1}{2}$

5.  
  
\_\_\_\_\_

6. The tide at a boat dock can be modeled by the equation  $y = -2 \cos\left(\frac{\pi}{6}t\right) + 8$ , where  $t$  is the number of hours past noon and  $y$  is the height of the tide, in feet. For how many hours between  $t = 0$  and  $t = 12$  is the tide at least 7 feet? [ The use of the accompanying grid is optional. ]

